

RESEARCH

Open Access



Magnitude and factors associated with iron supplementation among pregnant women in Southern and Eastern Regions of Ethiopia: Further Analysis of mini demographic and health survey 2019

Girma Teferi Mengistu^{1*}, Bizunesh Kefale Mengistu², Tolesa Gemeda Gudeta¹, Ayana Benti Terefe¹, Fedhesa Mamo Habtewold³, Mebratu Demissie Senbeta⁴, Seboka Abebe Sori⁴ and Hirut Dinku Jiru⁴

Abstract

Background: Anemia is a global public health problem that affects pregnant women. The most common cause of anemia is iron deficiency which is extremely common in developing countries. World health organization reported that 36.5% of pregnant women are anemic globally. In Ethiopia, 27.08% of women of the reproductive age group are anemic. Therefore, this study aimed to identify the magnitude and factors associated with iron supplementation during pregnancy in the southern and eastern regions of Ethiopia.

Methods: The data used in this analysis were extracted from Mini Demographic and Health Survey 2019. The survey was conducted in 9 regional states and two city administrations. The data used in the analysis were extracted from individual women datasets, and 1780 study participants were included in this study. The logistic regression analysis including bivariate and multivariable logistic regression at a 95% confidence interval and a *p*-value less than 0.05 was used.

Result: The finding of the study shows that iron supplementation during pregnancy in Southern and Eastern parts of Ethiopia was 50.06%. Among those who received iron, only about 20% took it for 90 days and more during their pregnancy. Iron supplementation among the pregnant women was affected by secondary education [AOR = 2.20, 95%CI (1.325, 3.638)], residing in urban [AOR = 1.75, 95%CI (1.192, 2.574)], having media at home [AOR = 1.41, 95%CI (1.022, 1.946)], having antenatal care follow up [AOR = 9.27, 95%CI (4.727, 18.169)], having 4 and more ANC follow up [AOR = 2.01, 95%CI (1.468, 2.760)], having antenatal care follow up at governmental health institutions [AOR = 3.40, 95%CI (1.934, 5.982)], and giving birth at governmental health institutions [AOR = 1.70, 95%CI (1.236, 2.336)].

Conclusion: Only one in two pregnant women was supplemented with iron during their recent pregnancy. The supplementation was affected by women's education, place of residence, presence of media at home, antenatal care follow-up, the number of antenatal care follow up, antenatal care follows up at governmental health institutions, and

*Correspondence: girmateferi56@gmail.com

¹ Department of Nursing, College of Medicine and Health Science, Wolkite University, Wolkite, Ethiopia

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

giving birth at the governmental health institution. The availability and accessibility of maternal care services and their functionality in providing maternal care services improve the supplementation.

Keywords: Pregnant women, Iron, Iron supplementation, Southern and eastern regions, Ethiopia

Introduction

Anemia in pregnancy is defined as a hemoglobin concentration of 11 gm/dl or less in the peripheral blood, which is diagnosed by examining characteristics of red blood cell changes in peripheral blood smear [1, 2]. Anemia occurs when there are not enough red blood cells or hemoglobin in the blood. Hemoglobin, which is the part of the red blood cells that carries oxygen, is made of iron [3]. Pregnant women are vulnerable to iron deficiency due to more iron being needed primarily to supply the growing fetus and placenta, which increases the maternal red cell mass [4]. The problem is that if pregnant women become deficient in these nutrients, they are not able to provide them an adequate amount for their baby, which can lead to anemia and increases the risk of complications [5].

Iron supplementation is the best way of reducing iron deficiency anemia [6]. Pregnant mothers are advised to take iron supplementation for 180 days during pregnancy [7]. In addition to iron supplementation, pregnant women should be advised to eat foods high in iron and prevent intestinal worms [8].

Anemia is a global public health issue that affects pregnant mothers [9]. Anemia during pregnancy results in a variety of life-threatening complications and poor pregnancy outcomes [3]. Iron deficiency is the most common dietary deficiency globally with negative maternal and perinatal outcomes [10]. It is extremely common in developing countries as a global epidemic where many women go through the whole pregnancy without reaching the required minimal iron intake [11]. Maternal anemia during the first or second trimester of pregnancy increases the risk of prematurity and being underweight at birth [12]. Due to increased iron requirements during pregnancy, iron deficiency can cause maternal anemia and reduce the iron reserves of the newborn [13]. Severe anemia can significantly increase the risk of maternal mortality and negatively affects fetal development [14]. But early initiation and use of a greater number of iron supplements reduce the risk of early neonatal complications [15].

According to World Health Organization estimates in 2019, the prevalence of anemia among pregnant women was 36.5% globally [16]. Similarly, the finding of a study conducted among 10 east African countries shows that the prevalence of anemia among reproductive-aged women was 34.85%. This study identified

that the prevalence ranges from 19.23% in Rwanda to 53.98% in Mozambique [17]. The report from further analysis of Ethiopia demographic and health survey (EDHS) 2016 shows that 27.08% of reproductive-aged women were anemic in Ethiopia [18]. It is highest in the Somali and Afar regions with 68.3% and 47.2% respectively [19].

Even though iron supplementation is used to prevent iron deficiency anemia in pregnancy, the report of EDHS 2016 shows that iron supplementation during pregnancy was 42% in Ethiopia [20]. The supplementation was showing improvement as evidenced by the report of Ethiopia Mini Demographic and Health Survey (EMDHS) 2019. This report identified that iron supplementation during pregnancy was 60% [8].

Iron supplementation during pregnancy can reduce maternal anemia at term by 70% by increasing their hemoglobin concentration [5]. However, it is affected by the women's level of adherence to the supplementation. The finding from the systematic review done on adherence to iron and folic acid supplementation in Ethiopia shows that nationally only 46.15% of the women adhered during pregnancy [21].

There are plenty of studies that have been done to identify factors associated with compliance to iron supplementation during pregnancy [22–26]. These studies do not address iron factors associated with supplementation. There was one study reporting determinants of iron supplementation nationally which reported a low level of iron supplementation in southern and eastern regions of the country [27]. Despite the low level of iron supplementation in these regions, there was a lack of evidence reporting factors affecting iron supplementation in the regions during pregnancy.

Therefore, this study aimed to identify the magnitude and factors associated with iron supplementation in the Southern and Eastern Regions of Ethiopia based on the Ethiopia mini demographic and health survey 2019. The finding of the study will help in improving maternal healthcare services.

Methodology

Data source

The data used in this analysis were obtained from Ethiopia Mini Demographic and Health Survey (DHS) 2019. Mini DHS 2019 is a national-level survey that gathers information about children, women, and men and

analyses and interprets the findings. The survey was done nationally among 9 Regional States and two City Administrations found in the country. The data collection period for mini DHS 2019 was from March and June 2019 [8].

Sampling procedure

This survey used two stages of sampling frame to collect the data at the national level. During the first stage, all census enumeration areas (EAs) created for the 2019 Population and Housing Census were used. And 305 EAs were selected with probability proportional to EA size and with independent selection in each sampling stratum. The second stage of selection used a fixed number of 30 households per cluster with an equal probability of systematic selection from the newly created household list. All women of reproductive age, fulfilling the selection criteria were eligible for the survey and selected for interviews [8].

For the current study, we used the 4 Regional States and one City Administration found in the Eastern and Southern parts of the country. The regions included in this analysis were Southern Nation Nationalities and Peoples Region, Harari Region, Dire Dawa City Administration, Somali Region, and Afar Region. This study used data from individual women datasets and a total weighted sample of 1780 women of reproductive age was included in the study. This sample was extracted from 8885 women interviewed during the survey. Among the respondents who participated in the survey, the data of 3964 respondents were recorded for the outcome variable at the national level. Then, since this study was based on the data from southern and eastern regions of the country, we extracted 1780 samples based on the regions included in the analysis.

Variables

Dependent

The dependent variable for the study was factors associated with iron supplementation during pregnancy. The variable was derived from the individual women dataset which has maternal-related information. The outcome variable was categorized and coded as 'Yes' (1) and 'No' (0).

Independent

The independent variables included in the current study were categorized under two categories. The first category was sociodemographic variables such as age (' ≤ 24 ', '25–29', '30–34', ' ≥ 35 '), educational attainment ('no education', 'primary education', 'secondary education and above'), media in a home ('yes', 'no', whether television or radio is present in the home), place of

residence ('urban', 'rural'), Region ('Afar', 'Somali', 'SNNPR', 'Harari', 'Dire Dawa'), marital status ('unmarried/in a relationship', 'married', 'widowed/divorced/separated'), wealth index ('poor', 'middle', 'rich'). The second category was obstetric characteristics: pregnant during the survey ('yes', 'no'), has ANC follow up in the last 5 years preceding the survey ('yes', 'no'), number of ANC follow ups for the last recent pregnancy in the last 5 years preceding the survey ('less than', '4 more'), place of ANC follow up ('home', 'governmental health institutions', 'private health facilities'), age of respondent at first birth (' ≤ 14 ', '15–19', '20–24', ' ≥ 25 '), place of delivery ('home', 'governmental health institutions', 'private health institutions'), delivery by cesarean section ('yes', 'no'), number of births in last 5 years ('1–2', ' ≥ 3 '), number of births in last 3 years ('no births', '1 birth', '2–3 births'), and number of living children ('no children', '1–2', '3–4', ' ≥ 5 ').

Operational definition

Iron supplementation during pregnancy: If the respondents were supplemented or told to buy iron tablets or syrup from the pharmacy during their antenatal care follow-up.

Data analysis

The extracted data were analyzed using computer software, SPSS version 23. The analysis of the data included both descriptive and inferential statistics. The descriptive analysis was used to analyze the frequency distribution of the data. The descriptive analysis included a presentation of data using frequency tables and figures with their description. The logistic regression analyses were used to check for the relation of the independent variable with the dependent variable, iron supplementation during pregnancy. Multicollinearity was checked for independent variables used in multivariable logistic regression analysis using the variation inflation factor. The logistic regression included bivariate and multivariable logistic regression analysis at a 95% confidence interval and p -value less than 0.05. In bivariate logistic regression, each variable was checked with the outcome variable. Those variables with a p -value of less than 0.25 were used for multivariable logistic regression analysis. Then variables with a p -value of less than 0.05 in multivariable logistic regression were declared as statistically significant with iron supplementation during pregnancy. The output of the logistic regression analysis was presented with an expression of odd ratios. The crude odds ratio was used for bivariate logistic regression while the adjusted odds ratio was used for multivariable logistic regression.

Ethical consideration

The authors received permission to use the data from DHS, and the data set is available online at <http://www.dhsprogram.com/data/available-datasets.cfm>.

Results

Sociodemographic characteristics

There were 1780 samples included in the analysis of the study. Among the study participants, 576 (32.4%) were found in the age category of 25–29. Regarding the educational status of the women 1065 (59.8%) of them were not educated, followed by primary education 503 (28.3%). Nearly two-thirds, 1128 (63.4%) of the respondents reported that they have media in their homes. The finding of the study also indicated that almost three fourth 1313 (73.8%) of them were from rural areas. Regarding marital status, almost all 1671 (93.9%) of the respondents were married. Almost one-half, 942 (52.9%) of the respondents were from poor families (Table 1).

Obstetrics characteristics

According to the current study, ANC service utilization was 1128 (63.4%), where only 575 (32.3%) of them had completed 4 visits (Fig. 1). Among those who had ANC follow up 981 (55.1%) of them were at governmental

health institutions. Regarding the age of respondents at first birth, about half 852 (47.8%) of them gave birth between the age category of 15–19, while 258 (14.5%) of the gave birth before the age of 14 years old. More than one-half, 958 (53.8%) of the mothers gave birth at home (Fig. 2) as evidenced by the finding of the study (Table 2).

Iron supplementation

The finding from the analysis of the data shows that iron supplementation during pregnancy in Southern and Eastern parts of Ethiopia was 50.06% (Fig. 3). Among those who received iron only about 20% took it for 90 days and more during their pregnancy (Fig. 4).


Factors associated with iron supplementation during pregnancy

The logistic regression analysis was conducted to identify the relation between the dependent and independent variables. We conducted bivariate and multivariable logistic regression analysis at 95%CI and declared a significant association with a *p*-value < 5%.


The bivariate analysis indicated that age of respondents, educational status, residence, region, media at home, ANC follow up, place of ANC follow up,

Table 1 Sociodemographic characteristics of reproductive age women, southern and eastern regions of Ethiopia

Variables	Characteristics	Frequency	Percent
Age of respondent	≤ 24	464	26.1
	25–29	576	32.4
	30–34	360	20.2
	≥ 35	380	21.3
Educational attainment	No education	1065	59.8
	Primary education	503	28.3
	Secondary and above	212	11.9
Media in home	Yes	1128	63.4
	No	652	36.6
Place of residence	Urban	467	26.2
	Rural	1313	73.8
Region	Afar	387	21.7
	Somali	340	19.1
	SNNPR	466	26.2
	Harari	307	17.2
	Dire Dawa	280	15.7
Marital status	unmarried/in relationship	15	0.8
	Married	1671	93.9
	widowed/divorced/separated	94	5.3
Wealth index	Poor	942	52.9
	Middle	177	9.9
	Rich	661	37.1



number of ANC follow up, place of delivery, number of births in the last 5 years, the number of births in the last 3 years, the number of living children were statistically significant with the outcome variable (iron supplementation). After controlling the confounding factors in Multivariable analysis, educational status, media in a home, residence, region, ANC follow up,



number of ANC follow up, ANC visit at governmental health institutions, and place of delivery was statistically significant.

Women with secondary education and above had 2.20 [AOR = 2.20, 95%CI (1.325, 3.638)] times higher odds of iron supplementation during pregnancy than women with no education. Mothers who live in urban areas had 1.75 [AOR = 1.75, 95%CI (1.192, 2.574)] times higher odds of iron supplementation during pregnancy than mothers living in rural areas. Women who had a television or radio in their home had 1.41 [AOR = 1.41, 95%CI (1.022, 1.946)] times higher odds of iron supplementation than mothers who had no television or radio in their home. Mothers who had ANC follow-up had 9.27 [AOR = 9.27, 95%CI (4.727, 18.169)] times higher odds of iron supplementation than mothers who had no ANC follow-up. Mothers who have four and more ANC follow-ups had 2.01 [AOR = 2.01, 95%CI (1.468, 2.760)] times higher odds of iron supplementation than mothers who had less than 4 ANC follow up. Mothers who had ANC follow-up at government health institutions had 3.40 [AOR = 3.40, 95%CI (1.934, 5.982)] times higher odds of iron supplementation than mothers who had no ANC follow-up at governmental health institutions. Women who gave birth at governmental health institutions had 1.70 [AOR = 1.70, 95%CI (1.236, 2.336)] times higher odds of iron supplementation than mothers who gave birth at home (Table 3).

Discussion

The finding of the current study identified that the magnitude of iron supplementation during pregnancy was 50.06% in the Southern and Eastern Regions of the country. This finding was lower than studies conducted in Germany, 65.2% [28], Khartoum Sudan, 92.1% [29], and Ethiopia, 60.87% [27]. The inconsistency could be related to geographical and socio-economical differences. Regions with poor socioeconomic status and remote areas have barriers to health care service [30].

On the other hand, this finding was higher than studies reported from Tanzania, 22.3% [31], and eight rural districts of Ethiopia, 35.4% [22]. These two studies were done before 14 years and 8 years respectively. The health delivery system is improving with time frame even though it is not satisfactory in some of the regions.

Among those who received iron supplementation, only 20% of them took it for 90 days and above during their pregnancy which is supported by another study [27]. On the other hand, the supplementation has increased when compared to the EDHS 2016 report where 42% of

Table 2 Obstetrics characteristics of reproductive age women, southern and eastern regions of Ethiopia

Variables	Characteristics	Frequency	Percent
Pregnant during survey	No	1516	85.2
	Yes	264	14.8
ANC follow up	No	652	36.6
	Yes	1128	63.4
Number of ANC follow up	Less than 4	1205	67.7
	4 and more	575	32.3
ANC follows up at home	No	1765	99.2
	Yes	15	.8
ANC follows up at Government His	No	799	44.9
	Yes	981	55.1
ANC follows up at Private/NGO His	No	1630	91.6
	Yes	150	8.4
Age of respondent at first birth	≤ 14	258	14.5
	15–19	851	47.8
	20–24	521	29.3
	≥ 25 +	150	8.4
Place of delivery	Home	958	53.8
	Governmental health institution	742	41.7
	Private health facility	80	4.5
Delivery by cesarean section	No	1662	93.4
	Yes	118	6.6
Number of births in last 5 years	1–2	1622	91.1
	≥ 3	158	8.9
Number of births in last 3 years	No birth	394	22.1
	1 birth	1143	64.2
	2–3 births	243	13.7
Number of living children	no child	22	1.2
	1–2	681	38.3
	3–4	494	27.8
	≥ 5	583	32.8





Fig. 3 Iron supplementation in Southern and Eastern Regions of Ethiopia

the mother were supplemented with iron during pregnancy [20]. Regionally the supplementation was lowest in the Somali Region where less than one-fifth of pregnant mothers received the supplements. Somali and Afar regions had a higher prevalence of anemia, 68.3% and 47.2% respectively [19]. But women of reproductive age have perceived barriers to healthcare service access, 69.9% [30].

The percentage of pregnant women who received antenatal care service was 63.4%, and only 32.3% of them has 4 or more ANC visits. This finding is similar to the study reported five years back, 62% of pregnant women received ANC service as evidenced by the report of EDHS 2016 [20]. On the other hand, the EMDHS 2019 report shows that the utilization of ANC service was 74.4%, which is higher than the current finding [27]. The difference between the studies indicates that the health



delivery system of these Regions is fragile even affecting the report at the country level [27].

In those Regions, institutional delivery was less than half, where only 46.18% of the pregnant mothers gave birth at governmental or private health institutions. More than half of the mothers gave birth at home without the assistance of health professionals. The mothers gave birth at home due to a lack of awareness about maternal care benefits [32]. Institutional delivery was slightly higher at the national level, 52.5% as reported by EMDHS 2019. Additionally, EDHS 2016 report shows that 48% of pregnant women gave birth, five years preceding the survey [20]. The service provision in these Regions indicates that it is five years backward from the other parts of the country. The difference in the report could be related to the geographical location and access to facilities such availability of health institutions and roads. The finding of the study identified that 14.5% of mothers gave birth before the age of 14. Delivery before the age of 14 years old can result from early marriage caused by poor socioeconomic status, lack of education at the community level, and cultural or religious related [33].

Iron supplementation was determined by the educational status of mothers, having media at home, place of residence, Region, ANC follow-up, number of ANC follow up, and place of delivery.

Educational status has a positive effect on iron supplementations [34]. Mothers with secondary and above education had 2.20 higher odds of iron supplementation during pregnancy than mothers with no education. This

could be related to exposure to information about the purpose of iron during pregnancy. Mothers who live in urban had 1.75 higher odds of iron supplementation during pregnancy than mothers from rural areas. This could have resulted from access to health facilities and information related to iron supplementation. Those mothers who had a radio or television in their home had 1.41 higher odds of iron supplementation during pregnancy than those who had no radio or television where they could acquire information.

Mothers living in the Somali region had lower odds of iron supplementation during pregnancy than mothers living in the SNNP region. This finding reflects the health delivery system of the region is not coordinated. On the other hand, mothers living in Afar and Harari had higher odds of receiving iron supplementations than mothers living in SNNPR. The population living in Afar and Harari regions is lower than SNNPR, and they could afford the health care delivery system.

Mothers who had ANC follow-ups had 9.27 higher odds of iron supplementation during pregnancy than those who had no ANC follow up. Mothers who had 4 or more ANC follow-ups had 2.01 higher odds of iron supplementation during pregnancy than those who had less than 4 ANC follow up. It is supported by a study from Sudan [29]. During ANC visits, iron supplementation is provided as one component of ANC service. Those mothers who had ANC follow-up at governmental health institutions had 3.40 higher odds of iron supplementation during pregnancy than others which is supported by another study [35]. At governmental health institutions, iron is freely supplemented. Mothers who gave birth at governmental health institutions had 1.70 higher odds of iron supplementation during pregnancy than those who gave birth at home. Mothers are counseled after delivery about their health and nutrition. Depending on their health status, they could be informed about the prevention of anemia through iron supplementation.

This study was done based on secondary data which was extracted from EMDHS 2019, and four regions and one city administration found in the southern and eastern parts of the country were studied. In this study, variables were excluded from the analysis because of unregistered data or missing values. Among the variables, hemoglobin and level of anemia were not included since no data was entered for the variables.

Conclusion

Even though iron supplementation during pregnancy prevents iron deficiency anemia, the current study identified that only one in two were supplemented. It

Table 3 Determinants of iron supplementation for pregnant women, Southern and Eastern Regions of Ethiopia

Variable	Characteristics	Received Iron		COR95%CI	AOR95%CI
		Yes	No		
Age of respondent	≤ 24	240(51.7%)	224(48.3%)	1.52(1.158, 2.001)**	
	25–29	311(54.0%)	265(46.0%)	1.66(1.283, 2.166)***	
	30–34	183(50.8%)	177(49.2%)	1.46(1.098, 1.964)*	
	≥ 35	157(41.3%)	223(58.7%)	1	
Educational attainment	No education	409(38.4%)	656(61.6%)	1	1
	Primary education	304(60.4%)	199(39.6%)	2.45(1.972, 3.045)***	1.21(0.883, 1.648)
	Secondary & above	178(84.0%)	34(16.0%)	8.39(5.702, 12.366)***	1.19(0.872, 1.630)**
Residence	Urban	331(70.9%)	136(29.1%)	3.27(2.606, 4.109)***	2.20 (1.325, 3.638)**
	Rural	560(42.7%)	753(57.3%)	1	1
Media in home	No	421(64.6%)	231(35.4%)	1	1
	Yes	470(41.7%)	658(58.3%)	0.39(0.321, 0.478)***	1.41(1.022, 1.946)*
Region	SNNPR	190(49.1%)	197(50.9%)	1	1
	Somali	58(17.1%)	282(82.9%)	0.87(0.670, 1.149)	0.39(0.238, 0.626)***
	Afar	244(52.4%)	222(47.6%)	0.18(0.134, 0.262)***	0.68(0.460, 1.004)
	Harari	207(67.4%)	100(32.6%)	1.88(1.395, 2.542)***	1.17(0.740, 1.867)
	Dire Dawa	192(68.6%)	88(31.4%)	1.98(1.455, 2.709)***	0.93(0.582, 1.479)
ANC follow up	No	858(76.1%)	270(23.9%)	1	1
	Yes	33(5.1%)	619(94.9%)	59.60(40.930, 86.808)***	9.27(4.727, 18.169)***
Number of ANC follow up	Less than four	411(34.1%)	794(65.9%)	1	1
	Four and more	480(83.5%)	95(16.5%)	9.76(7.600, 12.537)***	2.01(1.468, 2.760)***
ANC follow up: Governmental HIs	No	131(16.4%)	668(83.6%)	1	1
	Yes	760(77.5%)	221(22.5%)	17.53(13.797, 22.288)***	3.40(1.934, 5.982)***
Place of delivery	Home	280(29.2%)	678(70.8%)	1	1
	Governmental HIs	563(75.9%)	179(24.1%)	7.61(6.122, 9.474)***	1.70(1.235, 2.333)**
	Private/NGO HIs	48(60.0%)	32(40.0%)	3.63(2.274, 5.803)***	0.97(0.483, 1.949)
Number of births in last 5 years	1–2	832(51.3%)	790(48.7%)	1.76(1.262, 2.475)**	
	≥ 3	59(37.3%)	99(62.7%)	1	
Number of births in last 3 years	no births	198(50.3%)	196(49.7%)	1.32(0.963, 1.831)	
	1 birth	588(51.4%)	555(48.6%)	1.39(1.053, 1.841)*	
	2–3 births	105(43.2%)	138(56.8%)	1	
Number of living children	no children	8(36.4%)	14(63.6%)	1	
	1–2	405(59.5%)	276(40.5%)	2.56(1.063, 6.204)*	
	3–4	238(48.2%)	256(51.8%)	1.62(.671, 3.948)	
	≥ 5	240(41.2%)	343(58.8%)	1.22(.506, 2.964)	

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

was also evidenced that among those who have supplemented, only one-fifth of them received for 90 and more days. Iron supplementation during pregnancy was affected by maternal education, place of residence, having media at home, having antenatal care follow-up, having antenatal care follow-up at a governmental health institution, and place of delivery of the last recent baby. Availability and functionality of maternal care services will facilitate iron supplementation during pregnancy. Access to a health facility in nearby areas of their residential area helps in enabling the mothers to receive the service.

Acknowledgements

The authors would like to thank the measure DHS program for providing the data set used in the analysis.

Authors' contributions

G.T., T.G., A.B. Wrote the main manuscript text, and F.M., M.D., S.A., and H.D. prepared figures, and tables, G.T. and B.K. analyzed the data and wrote the output. All authors reviewed and approved the manuscript.

Funding

The study was not funded.

Availability of data and materials

The dataset used in this study is available on the hand of the corresponding author upon reasonable request.

Declarations

Ethics approval and consent to participate

All methods were carried out following relevant guidelines and regulations of the Demographic and Health Surveys (DHS) program. Permission was obtained from DHS data archivists after registration. The study used secondary data and consent to participate is not applicable.

Consent for publication

Not applicable.

Competing of interests

The authors declare that there is no conflict of interest.

Author details

¹Department of Nursing, College of Medicine and Health Science, Wolkite University, Wolkite, Ethiopia. ²Department of Statistics, College of Natural and Computational Science, Ambo University, Ambo, Ethiopia. ³Department of Midwifery, Jimma University Medical Center, Jimma, Ethiopia. ⁴Department of Midwifery, College of Medicine and Health Science, Wolkite University, Wolkite, Ethiopia.

Received: 8 April 2022 Accepted: 12 July 2022

Published online: 18 July 2022

References

- Chowdhury S, Rahman M, Moniruddin ABM. Anemia in pregnancy. *Med Today*. 2014;26(1):49–52.
- Sabina S, Iftequar S, Zaheer Z, Khan MM, Khan S. An overview of anemia in pregnancy. *J Innov Pharm Biol Sci*. 2015;2(2):144–51.
- Liyew AM, Tesema GA, Alamneh TS, Worku MG, Teshale AB, Alem AZ, et al. Prevalence and determinants of anemia among pregnant women in East Africa: A multi-level analysis of recent Demographic and Health Surveys. *PLoS One*. 2021;16(4):e0250560. <https://doi.org/10.1371/journal.pone.0250560>.
- Moshi FV, Millanzi WC, Mwampagatwa I. Factors Associated With Uptake of Iron Supplement During Pregnancy Among Women of Reproductive Age in Tanzania: an Analysis of Data From the 2015 to 2016 Tanzania Demographic and Health Survey and Malaria Indicators Survey. *Frontiers in Public Health*. 2021;9. Available from: <https://doi.org/10.3389/fpubh.2021.604058>
- Peña-Rosas JP, De-Regil LM, Garcia-Casal MN, Dowswell T. Daily oral iron supplementation during pregnancy. *Cochrane Database Syst Rev*. 2015;(7). <https://doi.org/10.1002/14651858.CD004736.pub5>
- Gebremichael TG, Haftu H, Gereziher TA. Time to start and adherence to iron-folate supplement for pregnant women in antenatal care follow up. *Northern Ethiopia Patient Prefer Adherence*. 2019;13:1057.
- Solomon Y, Sema A, Menberu T. Adherence and associated factors to iron and folic acid supplementation among pregnant women attending antenatal care in public hospitals of Dire Dawa, Eastern Ethiopia. *Eur J Midwifery*. 2021;5(August):1–7. <https://doi.org/10.18332/ejm/138595>.
- Ethiopian Public Health Institute (EPHI) [Ethiopia] and ICF. 2021. Ethiopia Mini Demographic and Health Survey 2019: Final Report. Rockville, Maryland, USA EPHI ICF. 2019;
- Melku M, Addis Z, Alem M, Enawgaw B. Prevalence and predictors of maternal anemia during pregnancy in gondar, northwest ethiopia: an institutional based cross-sectional study. *Maggio A, editor. Anemia*. 2014;2014:108593. <https://doi.org/10.1155/2014/108593>.
- Imdad A, Bhutta ZA. Routine iron/folate supplementation during pregnancy: effect on maternal anaemia and birth outcomes. *Paediatr Perinat Epidemiol*. 2012;26(Suppl 1):168–77.
- Abu-Ouf NM, Jan MM. The impact of maternal iron deficiency and iron deficiency anemia on child's health. *Saudi Med J*. 2015 Feb;36(2):146–9. Available from: <https://pubmed.ncbi.nlm.nih.gov/25719576>
- Haider BA, Olofin I, Wang M, Spiegelman D, Ezzati M, Fawzi WW. Anaemia, prenatal iron use, and risk of adverse pregnancy outcomes: systematic review and meta-analysis. *BMJ*. 2013;346:f3443.
- Loy SL, Lim LM, Chan S-Y, Tan PT, Chee YL, Quah PL, et al. Iron status and risk factors of iron deficiency among pregnant women in Singapore: a cross-sectional study. *BMC Public Health*. 2019;19(1):397. <https://doi.org/10.1186/s12889-019-6736-y>.
- Juul SE, Derman RJ, Auerbach M. Perinatal iron deficiency: implications for mothers and infants. *Neonatology*. 2019;115(3):269–74. <https://doi.org/10.1159/000495978>.
- Nisar BY, Dibley MJ. Earlier initiation and use of a greater number of iron-folic acid supplements during pregnancy prevents early neonatal deaths in Nepal and Pakistan. *PLoS One*. 2014;9(11):e112446. <https://doi.org/10.1371/journal.pone.0112446>.
- World Health Organization. WHO Global Anaemia estimates, 2021 Edition; Prevalence of anaemia in women; Accessed date 10 May 2022. Available from: https://www.who.int/data/gho/data/themes/topics/anaemia_in_women_and_children
- Teshale AB, Tesema GA, Worku MG, Yeshaw Y, Tessema ZT. Anemia and its associated factors among women of reproductive age in eastern Africa: a multilevel mixed-effects generalized linear model. *PLoS One*. 2020;15(9):e0238957. <https://doi.org/10.1371/journal.pone.0238957>.
- SiffirArgawu A, GobeboMekebo G, Bedane K, KanthMakarla R, Kefale B, Senbetu T, et al. Prevalence and Determinants of anaemia among reproductive-age women in Ethiopia: a nationally representative cross-sectional study. *J Pharm Res Int*. 2021;33(59B):687–98.
- Liyew AM, Teshale AB. Individual and community level factors associated with anemia among lactating mothers in Ethiopia using data from Ethiopian demographic and health survey, 2016; a multilevel analysis. *BMC Public Health*. 2020;20(1):1–11.
- Ethiopia FDR. Ethiopia Demographic and Health Survey 2016. Addis Ababa Ethiopia, and Rockville: CSA ICF Maryland, USA; 2016.
- Desta M, Kassie B, Chanie H, Mulugeta H, Yirga T, Temesgen H, et al. Adherence of iron and folic acid supplementation and determinants among pregnant women in Ethiopia: a systematic review and meta-analysis. *Reprod Health*. 2019;16(1):182. <https://doi.org/10.1186/s12978-019-0848-9>.
- Gebremedhin S, Samuel A, Mamo G, Moges T, Assefa T. Coverage, compliance and factors associated with utilization of iron supplementation during pregnancy in eight rural districts of Ethiopia: a cross-sectional study. *BMC Public Health*. 2014;14(1):607. <https://doi.org/10.1186/1471-2458-14-607>.
- Fite MB, Roba KT, Oljira L, Tura AK, Yadeta TA. Compliance with Iron and Folic Acid Supplementation (IFAS) and associated factors among pregnant women in Sub-Saharan Africa: A systematic review and meta-analysis. *PLoS One*. 2021;16(4):e0249789. <https://doi.org/10.1371/journal.pone.0249789>.
- Birhanu TM, Birarra MK, Mekonnen FA. Compliance to iron and folic acid supplementation in pregnancy, Northwest Ethiopia. *BMC Res Notes*. 2018;11(1):345. <https://doi.org/10.1186/s13104-018-3433-3>.
- AregaSadore A, AbebeGebretsadik L, AmanHussen M. Compliance with iron-folate supplement and associated factors among antenatal care attendant mothers in Misha District, South Ethiopia: community based cross-sectional study. *Factor-Litvak PR, editor. J Environ Public Health*. 2015;2015:781973. <https://doi.org/10.1155/2015/781973>.
- Getachew M, Abay M, Zelalem H, Gebremedhin T, Grum T, Bayray A. Magnitude and factors associated with adherence to Iron-folic acid supplementation among pregnant women in Eritrean refugee camps, northern Ethiopia. *BMC Pregnancy Childbirth*. 2018;18(1):83. <https://doi.org/10.1186/s12884-018-1716-2>.
- Tamirat KS, Kebede FB, Gonete TZ, Tessema GA, Tessema ZT. Geographical variations and determinants of iron and folic acid supplementation during pregnancy in Ethiopia: analysis of 2019 mini demographic and health survey. *BMC Pregnancy Childbirth*. 2022;22(1):1–11.
- Demuth IR, Martin A, Weissenborn A. Iron supplementation during pregnancy – a cross-sectional study undertaken in four German states. *BMC Pregnancy Childbirth*. 2018;18(1):491. <https://doi.org/10.1186/s12884-018-2130-5>.
- Abdullahi H, Gasim GI, Saeed A, Imam AM, Adam I. Antenatal iron and folic acid supplementation use by pregnant women in Khartoum. *Sudan BMC Res Notes*. 2014;7(1):1–4.
- Tamirat KS, Tessema ZT, Kebede FB. Factors associated with the perceived barriers of health care access among reproductive-age women in Ethiopia: a secondary data analysis of 2016 Ethiopian demographic and health survey. *BMC Health Serv Res*. 2020;20(1):691. <https://doi.org/10.1186/s12913-020-05485-y>.
- Ogundipe O, Hoyo C, Østbye T, Oneko O, Manongi R, Lie RT, et al. Factors associated with prenatal folic acid and iron supplementation among 21,889 pregnant women in Northern Tanzania: a cross-sectional hospital-based

study. *BMC Public Health*. 2012;12(1):481. <https://doi.org/10.1186/1471-2458-12-481>.

32. Ibrhim MA, Demissie M, Medhanyie AA, Worku A, Berhane Y. Reasons for low level of skilled birth attendance in Afar pastoralist community, North East Ethiopia: a qualitative exploration. *Pan Afr Med J*. 2018;30:51. Available from: <https://pubmed.ncbi.nlm.nih.gov/30197742>
33. Malhotra A, Malhotra PBA. The Causes, Consequences and Solutions to Causes, Consequences and Solutions to Forced Child Marriage Child Marriage Child Marriage in the Developing World in the Developing World. 2010.
34. Niang K, Faye A, Tine JAD, Diongue FB, Ndiaye B, Ndiaye MB, et al. Determinants of iron consumption among pregnant women in southern Senegal. *Open J Obstet Gynecol*. 2016;7(1):41–50.
35. Cesar JA, de Carvalho Dumith S, Chrestani MAD, Mendoza-Sassi RA. Iron supplementation among pregnant women: results from a population-based survey study. *Rev Bras Epidemiol*. 2013;16:729–36.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

