

RESEARCH

Open Access



Maternal employment status and child age are positive determinants of minimum dietary diversity among children aged 6–23 months in Sagnarigu municipality, Ghana: a cross-sectional study

Ambrose Atosona^{1*}, Jawahir Abukari Mohammed¹, Huzaifa Issahaku¹, Khadija Saani¹, Hammond Yaw Addae² and Fusta Azupogo³

Abstract

Background Intake of a diversified diet is key to the prevention of malnutrition among children as it results in improved intake of energy and micronutrients, which are deemed critical for better nutritional status of children. This study assessed minimum dietary diversity (MDD) and its determinants among children aged 6–23 months in the Sagnarigu Municipality of Ghana.

Methods This was an analytical cross-sectional study, carried out in the Sagnarigu Municipality, Ghana and involved 369 mother-child pairs selected through a systematic random sampling. A semi-structured questionnaire was used to obtain respondents' socio-demographic characteristics, feeding practices, nutritional knowledge and anthropometry. MDD was assessed using a repeated 24-hour dietary recall method. Chi-square/Fisher exact test and binary logistic regression analysis were used to determine the factors associated with MDD.

Results The study found that 24.9% of the children were between the ages of 6–8 months and 75.1% were between the ages of 9–23 months. About 64.2% of the children met the MDD. Children of mothers who were self-employed were approximately 2 times more likely to meet the MDD as compared to children of mothers who were unemployed [Adjusted Odd Ratio (AOR): 1.93, 95% CI (1.13–3.31), $P=0.017$]. Also, children aged 9–23 months were approximately 14 times more likely to meet MDD as compared to younger children aged 6–8 months [AOR: 13.98, 95% CI (7.54–25.91), $P<0.001$].

Conclusion Our study suggests that maternal empowerment may have positive effects on improving the MDD of infants and young children.

Keywords Minimum dietary diversity, Nutrient adequacy, Children, Ghana

*Correspondence:

Ambrose Atosona
aatosona@yahoo.com

¹Department of Nutritional Sciences, School of Allied Health Sciences, University for Development Studies, Tamale, Ghana

²Nursing and Midwifery Training College, Kpembe, Ghana

³Faculty of Agriculture, Department of Family and Consumer Sciences, Food and Consumer Sciences, University for Development Studies, Tamale, Ghana



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Childhood malnutrition, particularly chronic malnutrition (stunting) remains the most problematic public health problem globally [1]. Stunting affects 22% [2] of the global population, 28.6% [3] of children in Sub-Saharan Africa and 18% of children in Ghana [4]. Ghana's Northern and North-East Regions have significantly higher rates of stunting, affecting approximately 30% of infants and young children [4]. Stunting prevalence stands at 47.6% [5] in the Sagnarigu Municipality.

Malnutrition in children results in delays in physical and motor development, reduced intelligence, more behavioural problems, poor social skills, and increased susceptibility to diseases [6, 7]. Almost 45% of mortalities in children under the age of five are caused by undernutrition and these largely occur in nations with low and intermediate levels of income [1]. More than two-thirds of these undernutrition-related mortalities are linked to inappropriate feeding practices in the first two years of life [8]. The rapid growth during this period increases the children's nutrient requirements, making them more susceptible to malnutrition [9]. Hence, appropriate feeding practices during this period (appropriate, safe, adequate and frequent feeding) are critical for their optimal growth and development [10].

MDD, a diet quality indicator for children aged 6–23 months [11], is defined as the intake of at least five or more food groups from the recommended eight food groups: (1) Breast milk, (2) grains, roots and tubers, (3) legumes and nuts, (4) dairy products, (5) flesh foods, (6) eggs, (7) vitamin A rich fruits and vegetables and (8) other fruits and vegetables [12]. It is one of the several indicators developed by World Health Organization (WHO) to offer simple, accurate, and reliable metrics for evaluating infant and young child feeding practices (IYCF) at the population level [13].

Meeting the MDD is key to the prevention of malnutrition among children as it results in improved intake of energy and nutrients which are deemed critical for better nutritional status [14]. Yet, many parents, particularly those in the low-and middle-income countries struggle to provide diversified diets for their children [3]. Globally, less than a quarter of children aged 6 to 23 months meet the MDD score and feeding frequency [15]. In Sub-Saharan Africa and Ghana, only 25.1% [3] and 35.3% of children aged 6–23 months receive MDD [16] respectively. The lack of variety in children's diets in most Sub-Saharan Africa nations has been attributed to household food insecurity [17, 18]. Maternal education, nutritional knowledge, number of children, antenatal and postnatal care visits and household monthly income have been identified as key determinants of MDD among children aged 6 to 23 months in some settings [19–22]. Identifying context-specific factors influencing MDD is pivotal in

designing effective interventions. This tailored approach holds promise in mitigating childhood malnutrition, as it enables strategies that account for diverse local conditions and individual needs. By addressing these determinants, we can enhance the effectiveness of efforts aimed at improving children's nutrition and overall well-being. Despite MDD being a known major determinant of child nutritional status [14] and the fact that the prevalence rate of child malnutrition (stunting) is extremely high (47.6%) in the Sagnarigu Municipality of Ghana [5], MDD and its context specific determinants in the Sagnarigu Municipality are currently unknown. This study, therefore, aimed to assess MDD and its determinants among children aged 6–23 months in the Sagnarigu Municipality of Ghana.

Methods

Study design, area and population

An analytical cross-sectional study design was used in the present study. The study was conducted in Kalpohini, Kpalsi, Sagnarigu and Wurshe communities in the Sagnarigu municipality of Northern Ghana from August to October, 2022. Eligible participants were mother-child pairs with the children aged 6–23 months who consented to participate in the study. Participants who were not sound minded were excluded from the study.

Sampling

Multi-stage sampling method was used in this study. Four study communities were selected through simple random sampling. All the 79 communities under Sagnarigu Municipality were listed and assigned specific numbers written on pieces of paper, put in a basin, mixed, and picked one at a time with replacement till the number of communities were reached. Systematic random sampling was used to select the households, where the households were selected according to a random starting point but with a fixed interval. This interval was calculated by dividing the total number of households by the desired sample size. In households with two or more eligible children, only one child was selected through simple random sampling. The population proportion to size method was utilized to calculate the sample size for each of the communities.

Sample size

The sample size for this study was calculated using the Cochran's formula:

$$N = \frac{z^2 p(1-p)}{M.E^2} \quad [23]$$

Where:

N is the sample size.

z is confidence interval (95%) which gives a critical value of 1.96.

p is the estimated proportion of an attribute present in the population. Prevalence of minimum dietary diversity is 35.3% in northern Ghana [16].

M.E is the desired level of precision (5%=0.05).

Hence, $N=351$. A 5% contingency was considered to cover up incomplete questionnaires. Hence, $N=369$.

Study variables

Dependent variable

The dependent variable was MDD.

Independent variables

The independent variables were caregiver characteristics (age, marital status, ethnicity, religion, educational level, occupation, income level, birth interval, antenatal care visits, post-natal care visits, place of delivery, breastfeeding status, timely initiation of breastfeeding and nutritional knowledge) and child characteristics (age, sex and history of illness in the past 2 weeks).

Data collection

Pretesting of questionnaire

The questionnaire pretesting was done by administering the questionnaires to 37 (10% of total sample size) caregivers with children aged 6–23 months in the study district; these caregivers did not participate in the study. This allowed for the researchers to fine-tune the questions for clarity and comprehension by the participants.

Socio-demographic characteristics

The pretested semi-structured questionnaire was used to take information on socio-demographic characteristics of the mother including age, marital status, ethnicity, religion, educational level, occupation, income level and sex of child. Data collection was facilitated by two trained research assistants.

Health service utilization and obstetric characteristics

Participants' characteristics including antenatal care (ANC) visits, post-natal care (PNC) visits, place of delivery and birth interval were also documented.

Caregivers knowledge on dietary diversity and feeding practices

Caregivers' knowledge on dietary diversity and feeding practices was assessed using a questionnaire adapted from Solomon et al. [24]. The questionnaire consisted of ten knowledge questions. Each correct answer (yes) was assigned a score of 1, while any wrong answer was assigned a zero (0) score. Mothers who got a score of 7 and above out of the ten knowledge questions were deemed to have good knowledge while mothers who got

a score of less than 7 were deemed to have poor knowledge [24].

Dietary diversity

Data on MDD was collected using the WHO indicators for assessing IYCF practices [25]. A 24-hour dietary recall, repeated in 20% of random sub-sample [26], was used to obtain food intake information. The 24-hour dietary recall was conducted on two non-consecutive days, one weekday and one weekend [27–29]. The respondents were asked to recall all foods eaten and beverages taken by the children in the previous 24 h prior to the interview. The dietary diversity score (DDS) was assessed by assigning a score of 1 to a food group if the child ate any food item from the food group and a score of 0 if no food item from the food group was consumed. A total of 8 food groups [25] were considered in this study: Breast milk; grains, roots and tubers; legumes, nuts and seeds; dairy products; flesh foods, eggs; vitamin A-rich fruits and vegetables and other fruits and vegetables. Consequently, the minimum possible DDS score was 0 (no food group consumed) and the maximum possible DDS score was 8 (all food groups consumed). A child was classified as having achieved the MDD if he/she consumed at least 5 food groups out of the 8.

Anthropometry

Length (cm) of child was measured without footwear using an infantometer (Seca, Germany) and weight (kg) was measured without clothing using a digital weighing scale (Seca, Germany). The child's age, sex, and measurements of weight and length were used to calculate the following growth indicators: length-for-age (stunting), weight-for-age (underweight) and weight-for-length (wasting) in accordance with the WHO 2006 child growth reference [30]. The cut off point for stunting, wasting and underweight was $-2SD$ from the median of WHO child growth standard.

Data analysis

The Statistical Package for Social Science (SPSS version 22) was used to analyze the data. The results were reported with descriptive statistics including frequencies, percentages, means, and standard deviations. Chi-square/Fisher exact test was used for bivariate analysis. The factors with $p < 0.25$ [31] in the bivariate analysis were selected for multivariate logistic regression analysis to determine predictors of MDD. $P < 0.05$ (at two-tailed test) was considered significant.

Results

Socio-demographic characteristics

The study had a 100% response rate. Close to two-thirds of the mothers were within 20–30 years of age (62.6%).

Table 1 Socio-demographic characteristics

Variable	Category	Frequency	Percentage
Mother's age (in years)	< 20	21	5.7
	20–30	231	62.6
	31–40	111	30.1
	> 40	6	1.6
Religion	Christianity	18	4.9
	Islam	351	95.1
Maternal marital status	Single	1	0.3
	Married	367	99.5
	Separated	1	0.3
Number of children	< 5	247	66.4
	6–10	97	26.3
	≥ 11	25	6.8
Household monthly income	< Ghc500	329	89.1
	Ghc 500–999	25	6.8
	Ghc 1000–1500	10	2.7
	> Ghc1500	5	1.4
Ethnicity	Dagomba	333	90.2
	Gonja	12	3.3
	Others	24	6.5
Employment status	Self-employed	203	55.0
	Employed	23	6.2
	Unemployed	143	38.8
Mother's highest educational level completed	None	170	46.1
	Primary	65	17.6
	Middle/JHS	58	15.7
	SHS/vocational training	40	10.8
Age of child (in months)	6–8	92	24.9
	9–23	277	75.1
Sex of child	Male	191	51.8
	Female	178	48.2

Table 2 Feeding practices, maternal nutritional knowledge and nutritional status of children

Variable	Category	Frequency	Percentage
Currently breastfeeding	Yes	356	96.5
	No	13	3.5
Timely initiation of breastfeeding	Yes	246	66.7
	No	123	33.3
Nutritional knowledge of mother	Good	285	77
	Poor	86	23
Stunted	Yes	90	24.4
	No	279	75.6
Wasted	Yes	75	20.3
	No	294	79.7
Underweight	Yes	90	24.4
	No	279	75.6
Child history of illness in the past 2 weeks	Yes	121	32.8
	No	248	67.2

The predominant religion was Islam (95.1%), and nearly all mothers were married (99.5%). Also, the main ethnic group was Dagombas (90.2%), with the majority (89.1%) having a household monthly income of less than 500 Ghana cedis. Also, the majority (66.4%) of the mothers had less than 5 children. Regarding education, most of the mothers (46.1%) did not have any formal education. The majority of the children were males (51.8%) and aged 9–23 months (75.1%) (Table 1).

Feeding practices, maternal nutritional knowledge and nutritional status of children

In the present study, about 96.5% of the children were breastfed. About two-thirds (66.7%) of them started breastfeeding within the first hour of delivery. With regards to mother's knowledge on IYCF practices, the majority (77.0%) had good knowledge. About 32.8% of the children fell ill 2 weeks prior to the study. The prevalence of wasting, stunting and underweight were 20.3%, 24.4% and 24.4% respectively (Table 2). Of the total number of children, 237 (64.2%) met the MDD score (Fig. 1).

Health service utilization and obstetric characteristics

About two-thirds (66.9%) of the mothers delivered at the hospital. Almost all (99.5%) mothers attended antenatal care services, with the majority (92.1%) of them going for more than 4 visits. The number of mothers who attended postnatal care visits was 363 (98.4%). Also, about two-thirds of the mothers (64.5%) had a birth interval of more than 2 years (Table 3).

Factors associated with minimum dietary diversity practice

Significant determinants of MDD at the bivariate level included employment status ($p=0.119$), educational level ($p=0.153$), frequency of antenatal care visits ($p=0.027$), sex of child ($p=0.059$) and age of child ($p<0.001$) (Table 4).

In the multivariable binary logistic regression analysis, employment status and age of child were the predictor variables. Children of mothers who were self-employed were twice more likely to meet the MDD as compared to children of mothers who were unemployed [AOR: 1.93, 95% CI (1.13–3.31), $p=0.017$]. The children aged 9–23 months were about 14 times more likely to meet MDD as compared to younger children (6–8 months) [AOR: 13.98, 95% CI (7.54–25.91), $p<0.001$] (Table 5).

Discussion

The consumption of a well-diversified diet prevents childhood malnutrition as it improves the intake of nutrients and energy. This study investigated the prevalence of MDD and its predictors among children aged 6–23 months in the Sagnarigu Municipality of Ghana.




Fig. 1 The percentage of children who met the minimum dietary diversity (MDD)

Table 3 Health service utilization and obstetric characteristics

Variable	Category	Frequency	Percentage
Place of delivery	At home	90	24.4
	Health center	32	8.7
	Hospital	247	66.9
Antenatal care visits	Yes	367	99.5
	No	2	0.5
Frequency of antenatal care visits	1	4	1.1
	2–3	23	6.2
	≥ 4	340	92.1
	None	2	0.5
Post-natal care visits	Yes	363	98.4
	No	6	1.6
Birth interval	1 year	91	24.6
	2 years	40	10.8
	More than 2 years	238	64.5

The study revealed that about 64.2% of the children met the MDD score which was higher than the national average of 41% reported in the 2022 Ghana Demographic and Health Survey Report [4]. This is higher than the rate reported in a cross-sectional study in Ghana (35.3%) [16]. Similarly, the rate reported in the current study is also higher than the rates reported in Ethiopia [32] and Bangladesh [33], where the rates were found to be 12.6% and 28.7% respectively. On the other hand, the finding of the present study is comparable to that of Solomon et al. [24] and Sekartaji et al. [34], who reported rates of 59.6% and 63.2% respectively. This variation in results may partly be attributed to differences in methodologies [35, 36], population characteristics [35], nutrition knowledge of mothers [37], economic conditions [38], women's empowerment and food security [39].

The present study also revealed a general trend towards higher dietary diversity as children age. The children aged 9–23 months were approximately 14 times more likely to meet the MDD as compared to younger children (6–8 months). This finding is consistent with the findings of

several studies [32, 40–42]. It could be that children typically acquire a broader range of dietary preferences as they grow older, leading to increased diversity in their family meal choices. This might be a result of late introduction of complementary foods and when introduced on time, only milk or cereal foods like porridge are usually given, as some mothers have the perception that, the younger the child, the poorer the ability of the child to digest other foods such as beef, eggs, green leafy vegetables, carrots, and bananas [32, 43]. Also, teething and increased infections during infancy (6–8 months) lead to loss of appetite, resulting in low dietary intake and low dietary diversity [38]. This may explain the finding of the present and previous studies [43, 44]. Our finding emphasises the need for tailored nutritional interventions to address the specific needs of different age groups within the early childhood period. This could involve nutrition education on transitioning from exclusive breastfeeding to the introduction of solid foods and ensuring a diverse diet.

In the present study, children of mothers who were self-employed (traders, farmers, hairdresser etc.) were twice more likely to meet the MDD as compared to children of mothers who were unemployed. In conformity with the finding of the present study, studies by Issaka et al. [45] and Belay et al. [43] revealed that working mothers are more likely to feed their children diversified diets as compared to non-working (unemployed) mothers. This might be the case because mothers who work are more likely to have higher incomes which may translate to higher purchasing power, thereby improving household food security [46]. Increased household food security is associated with increased MDD among children in Ghana [47]. The stark difference in MDD achievement between children of self-employed mothers and those of unemployed mothers underscores the potential impact of economic stability and parental employment on child nutrition. Policies aimed at promoting and supporting

Table 4 Bivariate analysis of factors associated with minimum dietary diversity

Characteristic	Total population N = 369(%)	MDD UNMET = 35.8(%)	MDD MET = 64.2(%)	P-Value
Age of mother				
<20	21(100)	5(23.8)	16(76.2)	0.325
20–30	231(100)	90(39.0)	141(61.0)	
31–40	111(100)	36(32.4)	75(67.6)	
>40	6(100)	1(16.7)	5(83.3)	
Ethnicity				
Dagomba	333(100)	118(35.4)	215(64.6)	0.883
Gonja	12(100)	5(41.7)	7(58.3)	
Others	24(100)	9(37.5)	15(62.5)	
Religion				
Christianity	18(100)	6(33.3)	12(66.7)	0.825
Islam	351(100)	126(35.9)	225(64.1)	
Marital status				
Single	1(100)	0(0.0)	1(100.0)	1.000
Married	367(100)	132(36.0)	235(64.0)	
Separated	1(100)	0(0.0)	1(100.0)	
Employment status				
Self-employed	203(100)	66(32.5)	137(67.5)	0.119
Employed	23(100)	6(26.1)	17(73.9)	
Unemployed	143(100)	60(42.0)	83(58.0)	
Education				
None	170(100)	64(37.6)	106(62.4)	0.153
Primary	65(100)	19(29.2)	46(70.8)	
Middle/JHS	58(100)	16(27.6)	42(72.4)	
SHS/vocational training	40(100)	20(50.0)	20(50.0)	
Tertiary	36(100)	13(36.1)	23(63.9)	
Household income				
< GH¢ 500	329(100)	113(34.3)	216(65.7)	0.27
GH¢ 500–999	25(100)	15(60.0)	10(40.0)	
GH¢ 1000–1500	10(100)	2(20.0)	8(80.0)	
> GH¢ 1500	5(100)	2(40.0)	3(60.0)	
Period between birth				
1 year	91(100)	40(43.9)	51(56.1)	0.260
2 years	40(100)	13(32.5)	27(67.5)	
> 2	238(100)	79(33.2)	159(66.8)	
ANC attendance				
Yes	367(100)	131(35.7)	236(64.3)	1.000
No	2(100)	1(50.0)	1(50.0)	
Frequency of ANC visits				
Once	4(100)	3(75.0)	1(25.0)	0.027
2–3 times	23(100)	13(56.5)	1(25.0)	
≥4 times	341(100)	116(34.0)	225(66.0)	
PNC attendance				
Yes	363(100)	130(35.8)	233(64.2)	1.000
No	6(100)	2(33.3)	4(66.7)	
Sex of child				
Male	191(100)	77(40.3)	114(59.7)	0.059
Female	178(100)	55(30.9)	123(69.1)	
Age of child				
6–8	92(100)	72(78.3)	20(21.7)	< 0.001
9–23	277(100)	60(21.7)	217(78.3)	
Illness in the past 2 weeks				
Yes	121(100)	39(32.2)	82(67.8)	

Table 4 (continued)

Characteristic	Total population N= 369(%)	MDD UNMET= 35.8(%)	MDD MET= 64.2(%)	P-Value
No	248(100)	93(37.5)	155(62.5)	0.322
Knowledge of mother				
Poor knowledge	85(100)	32(37.6)	53(62.4)	
Good knowledge	284(100)	100(35.2)	184(64.8)	0.681
Early initiation of breastfeeding				
Met	247(100)	86(34.8)	161(65.2)	
Unmet	121(100)	45(37.2)	76(62.8)	0.655

Table 5 Multivariable analysis of factors associated with minimum dietary diversity

Characteristics	Unadjusted odds ratio (95% confidence interval)	P- value	Adjusted odds ratio (95% confidence interval)	P-value
Age (months)				
6–8	1		1	
9–23	13.02(7.35–23.07)	< 0.001	13.98 (7.54–25.91)	< 0.001
Employment status				
Unemployed	1	1		
Self-Employed	0.67(0.43–1.04)	0.073	1.930 (1.13–3.31)	0.017
Employed	0.49 (0.182–1.31)	0.155	3.496 (0.834–14.652)	0.087
Education				
None	1		1	
Primary	0.68 (0.369–1.27)	0.229	1.541 (0.74–3.20)	0.247
Middle/JHS	0.63 (0.328–1.21)	0.168	2.14 (0.97–4.71)	0.059
SHS/vocational	1.66 (0.0828–3.31)	0.154	0.854 (0.36–2.014)	0.719
Tertiary	0.94(0.443–1.98)	0.863	0.648 (0.21–1.98)	0.446
ANC attendance				
Once	1		1	
2–3 times	0.43 (0.04–4.82)	0.496	3.38 (0.22–50.92)	0.379
≥4 times	0.17 (0.02–1.67)	0.129	3.64 (0.03–45.27)	0.315
Sex of child				
Female	1		1	
Male	1.51(0.98–2.32)	0.060	1.07(0.64–1.80)	0.786

self-employment opportunities, or income generation activities for women could have a beneficial impact on child nutrition.

Furthermore, child's sex, maternal educational status, and number of ANC visits by mothers were not significantly correlated with MDD at the multivariate level in the present study. In line with this finding, a similar study by Amoah et al. [48] revealed no relationship between MDD and these three factors. Contrarily, Sema et al. [49], in a study that determined the MDD and related factors among children aged 6–23 months, failed to show a correlation between MDD and child's gender, maternal educational status and number of ANC visits by mothers. The mixed findings could, at least in part, be attributed to the methodological variances among the studies [35, 36].

This study is not without limitations. The cross-sectional nature of the study makes it difficult to establish causal associations as exposure and outcome were measured simultaneously. Also, being a recall and self-reported study, recall bias might have affected the exact estimation of minimum dietary diversity. Despite

these, the study has shed insight into MDD and related characteristics among children aged 6 to 23 months in Sagnarigu Municipality for the first time. It is recommended that longitudinal studies should be carried out to establish cause and effect relationship between the dependent and independent variables.

Conclusion

The level of MDD practice was high in the study area. Maternal employment status and age of child were the significant determinants of MDD. Mothers, especially those unemployed, should be encouraged to feed their children diversified diets with emphasis on the transition period from exclusive breastfeeding to complementary feeding so as to reduce the risks of malnutrition.

Abbreviations

MDD	Minimum dietary diversity
WHO	World Health Organization
ANC	Antenatal care
PNC	Post-natal care
IYCF	Infant and young child feeding
CM	Centimeters

KG Kilogram
 SPSS Statistical Package for Social Sciences
 AOR Adjusted Odds Ratio

Acknowledgements

We are thankful to all the participants who partook in the study.

Author contributions

AA, JAM, HI and KS designed the study and performed data collection and analysis. AA drafted the manuscript. HYA and FA reviewed the manuscript and provided input. All authors read and approved the manuscript for publication.

Funding

The authors did not receive funding for this study.

Data availability

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethical approval and consent to participate

Permission to conduct the study was obtained from the Committee on Human Research, Publications and Ethics (CHRPE/AP/674/22) at Kwame Nkrumah University of Science and Technology, Ghana. The study was carried out in accordance with the ethical principles of Declaration of Helsinki. Permission was obtained from leaders of the communities of study. Written informed consent of participants was also obtained. Participants were also assured of confidentiality of their responses. For mothers younger than 18 years, written informed consent was obtained from their parents. For infants and illiterate participants, written informed consent was obtained from their legal guardians/parents. Thumb-printed informed consent was obtained from the illiterate participants who were older than 16 years.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Received: 21 September 2023 / Accepted: 22 March 2024

Published online: 15 April 2024

References

- World Health Organization. Malnutrition. 2022 [cited 2022 November, 16]; Available from: <https://www.who.int/news-room/fact-sheets/detail/malnutrition>.
- Global Nutrition Report. The state of global nutrition. 2021 [cited 2022 15th November].
- Aboagye RG et al. Dietary diversity and undernutrition in children aged 6–23 months in Sub-Saharan Africa. *Nutrients*. 2021. 13(10).
- Ghana Statistical Service (GSS), Ghana Health Service (GHS), and ICF International, 2022 Ghana Demographic and Health Survey (DHS) Key Indicators Report. 2023, GSS: Rockville, Maryland, USA.
- Zorre EAK. *Curbing stunting growth In the Sagnerigu Municipality*. 2020 [cited 2022 November, 16]; Available from: <https://www.modernghana.com/news/1030311/curbing-stunting-growth-in-the-sagnerigu-municipal.html>.
- Skoet J, Stamoulis KG. The state of food insecurity in the world 2006: eradicating world hunger-taking stock ten years after the world food summit. *Food & Agriculture Org*; 2006.
- Black RE, Morris SS, Bryce J. Where and why are 10 million children dying every year? *Lancet*. 2003;361(9376):2226–34.
- World Health Organization. *Infant and young child feeding: model chapter for textbooks for medical students and allied health professionals*. World Health Organization: Geneva; 2009.
- Rao S, et al. Study of complementary feeding practices among mothers of children aged six months to two years-A study from coastal south India. *Australasian Med J*. 2011;4(5):252.
- Dwyer JT, et al. Feeding infants and toddlers Study 2008: progress, continuing concerns, and implications. *J Am Diet Assoc*. 2010;110(12):6.
- INDDEx Project. Minimum Dietary Diversity (MDD). 2022 [cited 2022 November, 17]; Available from: <https://inddex.nutrition.tufts.edu/inddex-project>.
- World Health Organization. *Global Nutrition Monitoring Framework: operational guidance for tracking progress in meeting targets for 2025*. Geneva: World Health Organization; 2017.
- World Health Organisation (WHO). *Indicators for assessing infant and young child feeding practices*. WHO: Geneva; 2008.
- Eshete T et al. *Determinants of inadequate minimum dietary diversity among children aged 6–23 months in Ethiopia: secondary data analysis from Ethiopian Demographic and Health Survey 2016*. *Agriculture & Food Security*, 2018. 7(1): p. 66.
- 2021 [cited 2022 World Health Organisation. *Infant and young child feeding*, November 17]. Available from: <https://www.who.int/news-room/fact-sheets/detail/infant-and-young-child-feeding>.
- Saaka M, et al. Magnitude and factors associated with appropriate complementary feeding among children 6–23 months in Northern Ghana. *BMC Nutr*. 2016;2(1):2.
- Dewey KG, Adu-Afarwah S. Systematic review of the efficacy and effectiveness of complementary feeding interventions in developing countries. *Matern Child Nutr*. 2008;4:24–85.
- Mekonnen TC, et al. Meal frequency and dietary diversity feeding practices among children 6–23 months of age in Wolaita Sodo town, Southern Ethiopia. *J Health Popul Nutr*. 2017;36(1):1–8.
- Edris MM, Atnafu NT, Abota TL. Determinants of dietary diversity score among children age between 6-23 months in bench maji zone southwest ethiopia. *Pediatr Health Res*. 2018;3(3):10.
- Temesgen H, Yeneabat T, Teshome M. Dietary diversity and associated factors among children aged 6–23 months in Sinan Woreda, Northwest Ethiopia: a cross-sectional study. *BMC Nutr*. 2018;4(1):5.
- Keno S, et al. Dietary diversity and associated factors among children aged 6 to 23 months in Chelia District, Ethiopia. *BMC Pediatr*. 2021;21(1):565.
- Belete KT, et al. Levels of dietary diversity and its associated factors among children aged 6–23 months in West Shoa, Ethiopia: a comparative cross-sectional study. *J Nutritional Sci*. 2022;11:e20.
- Uakarn C, Chaokromthong K, Sintao N. Sample size estimation using Yamane and Cochran and Krejcie and Morgan and Green formulas and Cohen Statistical Power Analysis by G*Power and comparisons. Volume 10. *APHEIT INTERNATIONAL JOURNAL*; 2021. pp. 76–88. 2.
- Solomon D, Aderaw Z, Tegegne TK. Minimum dietary diversity and associated factors among children aged 6–23 months in Addis Ababa, Ethiopia. *Int J Equity Health*. 2017;16(1):181.
- WHO/UNICEF. *Indicators for assessing infant and young child feeding practices: definitions and measurement methods*. Geneva; 2021.
- Tooze J. Estimating usual intakes from dietary surveys: Methodologic Challenges, Analysis Approaches, and recommendations for low- and Middle-Income Countries. Washington, DC: Intake— Center for Dietary Assessment/FHI Solutions; 2020.
- Agustina R, et al. Associations of meal patterning, dietary quality and diversity with anemia and overweight-obesity among Indonesian school-going adolescent girls in West Java. *PLoS ONE*. 2020;15(4):e0231519.
- National Cancer Institute. *The Dietary Assessment Primer-24-hour Dietary Recall (24HR) At a Glance*. 2024 [cited 2024 February 8]; Available from: <https://dietassessmentprimer.cancer.gov/profiles/recall/#:~:text=Collecting%20a%20recall%20for%20at,More%20about%20Usual%20Dietary%20Intake>.
- Cuenca-García M, et al. Association of breakfast consumption with objectively measured and self-reported physical activity, sedentary time and physical fitness in European adolescents: the HELENA (healthy lifestyle in Europe by Nutrition in Adolescence) Study. *Public Health Nutr*. 2014;17(10):2226–36.
- WHO. *Child growth standards*. 2023 [cited 2023 March 28]; Available from: <https://www.who.int/tools/child-growth-standards>.
- Bursac Z, et al. Purposeful selection of variables in logistic regression. *Source Code Biol Med*. 2008;3:17.
- Beyene M, Worku AG, Wassie MM. Dietary diversity, meal frequency and associated factors among infant and young children in Northwest Ethiopia: a cross-sectional study. *BMC Public Health*. 2015;15(1):1–9.
- Ahmed MS et al. *Association between dietary diversity and nutritional status of the children aged 6–23 months in Bangladesh: evidence from the Multiple Indicator Cluster Survey 2019* *Nutrire*, 2022. 47(1): p. 3.

34. Sekartaji R, et al. Dietary diversity and Associated factors among children aged 6–23 months in Indonesia. *J Pediatr Nurs*. 2021;56:30–4.
35. Kibret KT et al. *The effect of dietary patterns on maternal anaemia in North Shewa, Ethiopia: A case-control study with Propensity Score Analysis* *Nutrition and Health*, 2023: p. 02601060231152345.
36. Gebremichael MA, Lema TB, Diversity D. Nutritional status, and Associated factors among pregnant women in their first trimester of pregnancy in Ambo District, Western Ethiopia. *Nutr Metabolic Insights*. 2023;16:11786388231190515.
37. Opoku Agyemang G, et al. Factors associated with food consumption and dietary diversity among infants aged 6–18 months in Ashanti Region, Ghana. *PLoS ONE*. 2023;18(11):e0294864.
38. Merga G, et al. Dietary diversity and associated factors among women of reproductive age in Jeldu District, West Shoa Zone, Oromia Ethiopia. *PLoS ONE*. 2022;17(12):e0279223.
39. Lufuke M et al. Women's Empowerment, Food Security, and Nutrition Transition in Africa. *Int J Environ Res Public Health*, 2022. 20(1).
40. Bedada Damtie S, Benti Tefera T, Tegegne Haile M. *Dietary Diversity Practice and Associated Factors among Children Aged 6–23 Months in Robe Town, Bale Zone, Ethiopia* *J Nutr Metab*, 2020. 2020: p. 9190458.
41. Dangura D, Gebremedhin S. Dietary diversity and associated factors among children 6–23 months of age in Gorche district, Southern Ethiopia: cross-sectional study. *BMC Pediatr*. 2017;17(1):1–7.
42. Kuche D, et al. Factors associated with dietary diversity and length-for-age z-score in rural Ethiopian children aged 6–23 months: a novel approach to the analysis of baseline data from the sustainable undernutrition reduction in Ethiopia evaluation. *Matern Child Nutr*. 2020;16(1):e12852.
43. Belay DG, et al. Determinants of inadequate minimum dietary diversity intake among children aged 6–23 months in Sub-saharan Africa: Pooled Prevalence and Multilevel Analysis of Demographic and Health Survey in 33 sub-saharan African countries. *Front Nutr*. 2022;9:894552.
44. Mokori A, Schonfeldt H, Hendriks SL. Child factors associated with complementary feeding practices in Uganda. *South Afr J Clin Nutr*. 2017;30(1):7–14.
45. Issaka AI, et al. Determinants of suboptimal complementary feeding practices among children aged 6–23 months in seven francophone West African countries. *Matern Child Nutr*. 2015;11:31–52.
46. Antwi KD, Lyford CP. Socioeconomic determinants of rural households' food security status in Northern Ghana. *J Agric Food Sci*. 2021;19(2):86–94.
47. Antwi J et al. Household food insecurity is associated with child's dietary diversity score among primary school children in two districts in Ghana. *Food Nutr Res*, 2022. 66.
48. Amoah AN et al. Correlates of dietary diversity among children aged 6–23 months of head porters in Ghana. *Front Public Health*, 2022. 10.
49. Sema A, et al. Minimum Dietary Diversity Practice and Associated factors among children aged 6 to 23 months in Dire Dawa City, Eastern Ethiopia: A Community-based cross-sectional study. *Global Pediatr Health*. 2021;8:2333794X21996630.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.